

Dichoptic De-Masking Learning in Adults With Amblyopia and Its Mechanisms

Xiang-Yun Liu¹ and Jun-Yun Zhang²

¹Department of Ophthalmology, Tengzhou Central People's Hospital, Tengzhou, Shandong Province, China

²School of Psychological and Cognitive Sciences, and Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China

Correspondence: Jun-Yun Zhang, School of Psychological and Cognitive Sciences, and Beijing Key Laboratory of Behavior and Mental Health, Peking University, No. 5 Yiheyuan Road, Haidian District, Beijing 100871, China; zhangjunyun@pku.edu.cn.

Submitted: December 28, 2018

Accepted: May 15, 2019

Citation: Liu X-Y, Zhang J-Y. Dichoptic de-masking learning in adults with amblyopia and its mechanisms. *Invest Ophthalmol Vis Sci.* 2019;60:2968-2977. <https://doi.org/10.1167/iovs.18-26483>

PURPOSE. Recently, we reported that dichoptic de-masking training can further boost stereoacuity, but not visual acuity, in adults with amblyopia after extensive monocular perceptual training. Here, we investigated whether this dichoptic training targets on interocular suppression directly, or improves vision through high-level brain mechanisms.

METHODS. Eleven adults with amblyopia first used amblyopic eyes (AEs) to perform contrast ($n = 6$) or orientation ($n = 5$) discrimination training, while resisting dichoptic noise masking from fellow eyes (FEs). Learning was indicated by increased maximal tolerable noise contrast (TNC) for AE contrast/orientation discrimination. After dichoptic training, six observers continued to use AEs to perform monocular training for nine sessions.

RESULTS. (1) Training of dichoptic de-masking doubled maximal TNC, but learning did not transfer much to the same task at an orthogonal orientation or a different task, showing orientation/task specificities. (2) Following a training-plus-exposure (TPE) protocol, AEs then received exposure of the orthogonal orientation by performing the other orientation/contrast discrimination task at the orthogonal orientation. After this TPE training, dichoptic learning with the original discrimination task transferred to the orthogonal orientation. (3) Dichoptic training improved AE's acuity (1.2 lines), stereoacuity (60.2%), and contrast sensitivity (mainly at higher spatial frequencies). (4) Additional monocular training did not produce further acuity and stereoacuity gains.

CONCLUSIONS. The initial orientation/task specificities exclude the possibility that dichoptic training reduces physiological interocular suppression. The later transfer of learning to an orthogonal orientation with TPE training suggests improvement in high-level brain processing. Dichoptic training may strengthen top-down attention to AEs to counter the impacts of attentional bias to FEs and/or physiological interocular suppression and improve stereoacuity.

Keywords: amblyopia, dichoptic training, perceptual learning, orientation specificity, task specificity

Amblyopia is a developmental visual disorder due to abnormal binocular visual experience (e.g., strabismus and anisometropia) in early childhood that disrupts the development of the visual cortex.^{1,2} Imbalanced visual inputs from two eyes may lead to interocular suppression or inhibition of the amblyopic eye (AE) by the strong fellow eye (FE).³ As a consequence, visual acuity, stereoacuity, as well as many other visual functions, are compromised.^{4,5}

Many studies have demonstrated that perceptual learning improves vision in adults with amblyopia.^{6,7} Although amblyopia affects both binocular and monocular visual functions, earlier perceptual learning studies mostly perform monocular training in the AE with the FE patched. More recent studies employ dichoptic training, targeting abnormal binocular functions directly via reducing interocular suppression, strengthening binocular fusion, and promoting binocular vision. Many dichoptic training studies use signal integration training paradigms, in which the task elements are separated between the two eyes and must be integrated for successful task completion.⁸⁻¹⁵ Dichoptic training may assist information

integration from the two eyes to help recover stereovision in amblyopic patients.⁶

In a previous study, we adopted a different dichoptic de-masking training paradigm (details provided in Methods and Results sections), in which the observers were trained to discriminate the contrast or orientation of a Gabor stimulus presented to the AE while discounting the masking effect from a noise masker presented to the FE.¹⁶ Dichoptic de-masking training was performed by a group of monocularly well-trained adult amblyopic observers to isolate the effects of dichoptic training. The observers were significantly more capable of discounting dichoptic noise masking after training. Moreover, dichoptic training produced extra gains of stereoacuity, but not visual acuity, in these monocularly well-trained amblyopic observers, supporting Levi et al.⁶ on the potential advantages of dichoptic training.

Like in adults with normal vision, monocular perceptual learning in those with amblyopia is often specific to the trained orientation. The orientation specificity has been attributed to training induced neural plasticity in the amblyopic early visual





contrast, the high-level hypothesis supposes that dichoptic study followed the tenets of the Declaration of Helsinki and

was approved by the institutional review board of Peking University.

Study Design

The basic experimental design is represented schematically in Figure 1A. Prior to training the visual acuities and contrast sensitivity functions for both amblyopic and fellow eyes, as well as the stereoacuity, were measured. Eleven observers were assigned into two groups randomly. Following a dichoptic TPE protocol: (1) The first group ($n = 6$) practiced contrast discrimination at a vertical orientation for nine sessions. Then they received exposure to the orthogonal orientation through an irrelevant orientation discrimination task for five sessions. (2) The second group ($n = 5$) first practiced orientation discrimination at a horizontal orientation for five sessions. Then they received exposure to the orthogonal orientation through an irrelevant contrast discrimination task for another five sessions. After the dichoptic TPE training, the visual acuities, contrast sensitivity functions, and stereoacuity were remeasured. A subset of observers ($n = 6$; S1, S2, S3, S5, S7, and S11 in the Table) then performed monocular orientation training for nine sessions. After this monocular training the visual acuities and stereoacuity were remeasured.

Apparatus and Stimuli

The setup was identical to that in Liu and Zhang.¹⁶ The stimuli were generated with Psychtoolbox-3 software²⁷ and presented on a 21-in Sony G520 CRT monitor (2048×1536 pixel, 0.19×0.19 mm/pixel, and 75-Hz frame rate). The head of the observer was stabilized by a chin-and-head rest. Experiments were run in a dimly lit room. For grating acuity and contrast sensitivity testing, a 14-bit look-up table achieved with a video attenuator was used to linearize the luminance of the monitor (mean luminance = 27 cd/m^2), and for other tasks an 8-bit look-up table was used (mean luminance = 50 cd/m^2).

The dichoptic stimuli (Fig. 1B) consisted of a pair of collinear vertical or horizontal Gabors (Gaussian windowed sinusoidal gratings) presented in AE and a band-pass filtered white noise masker in FE. The two Gabors had the same spatial frequency at 40% of AE's cut-off frequency, standard deviation at 1 wavelength (the reciprocal of spatial frequency),

orientation at 0° or 90° , phase at 90° , and a center-to-center distance of 4 wavelengths. The cut-off frequency of AE (Mean = 14.4 cpd , SD = 3.6 cpd) was assessed by a grating acuity test for each observer before training. The viewing distance was 1.2 m. In contrast discrimination trials, one Gabor's contrast was set at 0.80, and the other Gabor's contrast was $0.80 - 1.414 \times$ contrast discrimination threshold (with no masker presented in FE). The contrast discrimination threshold was premeasured for each observer with the same Gabor stimulus at a reference contrast of 0.80 (AE's contrast just-noticeable difference (JND) threshold: mean = 0.189 , SD = 0.031). In orientation discrimination trials, the global orientation of two always aligned Gabors were tilted upper or lower from horizontal. The orientation offset was 1.414 times the orientation discrimination threshold premeasured for each observer with no masker presented in FE (AE's orientation JND threshold: mean = 1.5° , SD = 0.3°). The contrast of two Gabors was identical at 0.80.

The band-pass filtered noise masker was 512×512 pixels ($4.4^\circ \times 4.4^\circ$) in size. To create the noise masker, a 512×512 pixels zero-mean white noise field was first generated, with each element being 2×2 pixels. The white noise field was then filtered in the frequency domain by a 1-octave band-pass filter centered at the same frequency of the Gabors. A new noise masker was generated every trial.

The stimulus for monocular orientation discriminat

9.80-4578 -1

observer pressed the space bar to initiate the trial as soon as the whole cross appeared stable. Immediately after the key press, a black square contour ($1.5^\circ \times 1.5^\circ$, the contour lines were 2-arcmin thick) was presented for 200 ms to prime attention to AE. After that the Gabor stimuli and the noise masker were presented dichoptically for 200 ms.

In the contrast discrimination trials, the observers were asked to judge which Gabor had a higher contrast. In the orientation discrimination trials, they were asked whether the 2-Gabor stimuli tilted upper or lower from horizontal. A staircase varied the root mean square contrast of the noise masker upon AE's contrast or orientation judgment. The staircase followed a 3-up-1-down rule that resulted in a 79.4% convergence rate. Specifically, three consecutive correct responses would raise the noise contrast by one step, and one incorrect response would lower the noise contrast by one step. The step size of the staircase was 0.05 log units. Each staircase consisted of eight reversals (~40-50 trials). The geometric mean of the last six reversals was taken as the maximal tolerable noise contrast (TNC) for successful contrast or orientation discrimination.

To ensure effective noise masking (i.e., an observer did not close his/her fellow eye), in 20% of the trials a white digit ("1" or "2," $1.1^\circ \times 1.7^\circ$ in size) was centered on the noise masker in FE while a blank screen was presented in AE. The observer needed to report the digit by key press (the mean correct rate = $95.5 \pm 1.5\%$). Auditory feedback was given on incorrect responses in all trials.

The dichoptic TPE protocol consisted of a first training phase and a second exposure phase. Before and after the first training phase (i.e., contrast/orientation discrimination training), the following conditions were tested to evaluate the learning and transfer effects: (1) maximal TNC for AE's contrast/orientation discrimination at the trained orientation (groups 1, 2), and (2) maximal TNC for AE's contrast (group 1) or orientation discrimination (group 2) at an untrained orthogonal orientation. Each condition was measured for five staircases (~200-250 trials). After the second exposure phase (orientation/contrast discrimination training at an orthogonal orientation), only condition (2) was re-tested to evaluate the learning and transfer effects. All staircases were run following a randomly permuted table for each observer. The duration varied from 1 to 2 hours, depending on the conditions. In the training and exposure phases, each daily session consisted of 20 staircases (for a total number of 800~1000 trials) and lasted for approximately 2 hours. More details can be found in the Results section below.

During monocular training, orientation discrimination threshold was measured with a 2AFC staircase procedure in AE. In each trial, a foveal fixation cross was flashed for 400 ms before the onset of the stimulus. Then the reference and the test stimuli were presented separately in two 200-ms stimulus intervals in a random order, separated by a 500-ms interstimulus interval. Threshold was estimated following a 3-down-1-up staircase rule that resulted in a 79.4% convergence rate. The step size of the staircase was 0.05 log units. Each staircase consisted of two preliminary reversals and six experimental reversals. The geometric mean of the experimental reversals was taken as the threshold for each staircase run. Each session consisted of 20 staircases (for a total number of 800~1000 trials) and lasted for approximately 2 hours.

Interocular Suppression

Several studies have suggested that the interocular contrast ratio is a reliable objective measurement of interocular suppression.^{9,28} Therefore, we adopted the interocular contrast ratio, which was the maximal TNC for AE divided by the

maximal TNC for FE, to assess the strength of interocular suppression. Specifically, in the pre- and posttests, the Gabors and the noise masker were switched between eyes, so that the noise masker was presented to AE and the Gabor stimuli were presented to FE. Thus, the maximal TNCs for FE contrast

Stereoacuity. The stereoacuity was evaluated using the Randot Stereo Test (Stereo Optical Co., Inc., Chicago, IL, USA) with polarizing glasses at a 40-cm viewing distance under normal room lighting. The stereo test was administered and scored according to the manufacturer's instructions. A graded sequence test was provided by contoured circles at 10 levels of disparity ranging from 400 to 20 arcsec. Randot forms with disparities at 500 and 250 arcsec were also used to provide additional steps of disparity.

RESULTS

Dichoptic De-Masking Learning Showed Orientation and Task Specificity

Eleven adult amblyopic observers with no prior monocular training experience were randomly divided into two groups. The first group of six initially practiced AE contrast discrim-

ination at a vertical orientation with dichoptic noise masking for nine sessions (Fig. 2A). We used the percent improvement ($PI = [\text{threshold_post}/\text{threshold_pre} - 1] \times 100$) to index the learning and transfer effects. After the first training phase, the maximal TNC for AE contrast discrimination was significantly improved by $173.1 \pm 39.8\%$ ($t_5 = 4.35$, $P = 0.007$, Cohen's $d = 1.78$; 2-tailed paired t -test in this and later analyses unless

specificity in dichoptic de-masking learning. After initial contrast discrimination training, the amblyopic observers in group 1 continued to practice an AE orientation discrimination task with the same stimuli at an orthogonal orientation, also under dichoptic noise masking. The new orientation task alone had no impact on AE contrast discrimination because of the task specificity, but it exposed the observers to the orthogonal orientation.


Orientation discrimination by $69.5 \pm 11.7\%$ (Fig. 2A, $t_5 =$

$t_{10} = -1.24, P = 0.24$, Cohen's $d = 0.37$), indicating that training improved uncrowded acuity slightly more than crowded acuity.


Stereoacuity. The dichoptic training improved stereoacuity from $410.9'' \pm 70.7''$ to $152.7'' \pm 35.9''$, or by $60.2 \pm 4.9\%$ (Fig. 3D, $t_{10} = 9.04, P < 0.001$, Cohen's $d = 2.52$). Note that in Figure 3D, the mean and SD for S10 and S11 were initially $1000''$ and $1000''$ test disparity.

her stereoacuity to be 600 arcsec. The improvement of

Contrast Sensitivity



0.



51

...the trained orientation (Fig. 3.41).

...frequency at 23.0 ± 1.7 cpd ($P < 0.001$). After training, contrast sensitivity functions in AEs were improved but still showed contrast sensitivity loss, primarily at higher spatial



the binocular vision impairments caused by strabismic amblyopia. A recent hypothesis is that training leads to better attention to the AE, so as to ease the effects of direct interocular suppression in a top-down manner to improve vision.³³ In general, this hypothesis is consistent with our claim that perceptual learning in amblyopic observers, like in normals, is a high-level learning process,¹⁹ which may involve improved attention to the AE. In our dichoptic learning, the observers are purposely trained to counter the masking effects from the FE. Therefore, the improved attention to the AE would reduce the attentional bias to the FE, and/or counter-balance the low-level physiological interocular suppression in V1.²⁵ This would result in a lower interocular suppression index that may reflect both high-level attentional bias and low-level physiological interocular suppression, as shown in our data.

We understand that our current study has its limitations. First, it is possible that the results are specific to our particular dichoptic training paradigm. We present a masker in one eye and a target in the other eye. The training principles and underlying mechanisms may be distinct from other dichoptic training studies in which the task elements are separated between the two eyes and must be integrated for successful task completion.⁸⁻¹³ Second, our results are largely based on anisometric amblyopes (>70%). It is suggested that the mechanisms underlying strabismic and anisometric amblyopia are different.^{34,35} The applicability of our conclusions to other types of amblyopia needs to be experimented. Third, more observers need to be included to confirm the results that monocular training would bring no more benefits after dichoptic training.

In our study, six of 11 observers received new lenses, which they wore only during the training sessions for a total of 20 to 28 hours, while the other five wore their existing lenses. We found no significant difference of E-chart acuity improvements in AEs between these two subgroups of observers ($P = 0.08$). There are reports that for adults with amblyopia, refractive adaptation has limited and insignificant effects on visual acuity and stereoacuity.^{36,37} Therefore, we assume that the refractive adaptation effects from 20 to 28 hours of new lens wearing would have very small effects on acuity and stereoacuity improvements in these six observers, and the overall effects would be minimal when all 11 observers' results are considered together.

We did not perform follow-up measurements in the current study. However, follow-up measurements were carried out in a previous study of ours using the same training paradigm.¹⁶ In that study, seven of 13 amblyopic observers were retested 10 months (mean = 10.3 months, SD = 0.9 months) after they finished dichoptic training. The maximal tolerable noise contrasts were not significantly different from those measured immediately after training ($t_6 = 0.06$, $P = 0.96$, Cohen's $d = 0.03$). The stereoacuities were not significantly different either ($t_6 = 0$, $P > 0.99$, Cohen's $d = 0$). These results indicate that the dichoptic training effects can persist for an extended period.

CONCLUSIONS

We demonstrated that dichoptic de-masking learning of visual discrimination in adults with amblyopia can transfer nearly completely to an orthogonal orientation with a TPE protocol, and that the learning is task specific. These results suggest high-level dichoptic learning, in which the amblyopes may learn the rules of reading out orientation or contrast signals from dichoptically presented noise, so that learning is transferrable across orientations. Dichoptic training may improve top-down attention to the amblyopic eye, so as to

counter attentional bias to the FE and/or physiological interocular suppression.

Acknowledgments

The authors thank Cong Yu, Dennis Levi, and Lei Liu for their insightful comments and discussions.

Supported by Natural Science Foundation of China Grant 31470975 (JYZ; Beijing, China).

Disclosure: X.-Y. Liu, None; J.-Y. Zhang, None

References

- Birch EE. Amblyopia and binocular vision. *Prog Retin Eye Res.* 2013;33:67-84.
- Kiorpes L, Kiper DC, O'Keefe LP, Cavanaugh JR, Movshon JA. Neuronal correlates of amblyopia in the visual cortex of macaque monkeys with experimental strabismus and anisometropia. *J Neurosci.* 1998;18:6411-6424.
- Harrad R, Sengpiel F, Blakemore C. Physiology of suppression in strabismic amblyopia. *Br J Ophthalmol.* 1996;80:373-377.
- McKee SP, Levi DM, Movshon JA. The pattern of visual deficits in amblyopia. *J Vis.* 2003;3(5):380-405.
- Giaschi D, Lo R, Narasimhan S, Lyons C, Wilcox LM. Sparing of coarse stereopsis in stereodeficient children with a history of amblyopia. *J Vis.* 2013;13(10):17.
- Levi DM, Knill DC, Bavelier D. Stereopsis and amblyopia: a mini-review. *Vision Res.* 2015;114:17-30.
- Levi DM, Li RW. Perceptual learning as a potential treatment for amblyopia: a mini-review. *Vision Res.* 2009;49:2535-2549.
- Ding J, Levi DM. Recovery of stereopsis through perceptual learning in human adults with abnormal binocular vision. *Proc Natl Acad Sci U S A.* 2011;108:E733-E741.
- Hess RF, Mansouri B, Thompson B. A binocular approach to treating amblyopia: antisuppression therapy. *Optom Vis Sci.* 2010;87:697-704.
- Hess RF, Mansouri B, Thompson B. A new binocular approach to the treatment of amblyopia in adults well beyond the critical period of visual development. *Restor Neurol Neurosci.* 2010;28:793-802.
- Astle AT, McGraw PV, Webb BS. Recovery of stereo acuity in adults with amblyopia. *BMJ Case Rep.* 2011;2011:bcr0720103143.
- Ooi TL, Su YR, Natale DM, He ZJ. A push-pull treatment for strengthening the 'lazy eye' in amblyopia. *Curr Biol.* 2013;23:R309-R310.
- Vedamurthy I, Nahum M, Huang SJ, et al. A dichoptic custom-made action video game as a treatment for adult amblyopia. *Vision Res.* 2015;114:173-187.
- Xi J, Jia WL, Feng LX, Lu ZL, Huang CB. Perceptual learning improves stereoacuity in amblyopia. *Invest Ophthalmol Vis Sci.* 2014;55:2384-2391.
- Li J, Thompson B, Deng D, Chan LY, Yu M, Hess RF. Dichoptic training enables the adult amblyopic brain to learn. *Curr Biol.* 2013;23:R308-R309.
- Liu XY, Zhang JY. Dichoptic training in adults with amblyopia: additional stereoacuity gains over monocular training. *Vision Res.* 2018;152:84-90.
- Levi DM, Polat U. Neural plasticity in adults with amblyopia. *Proc Natl Acad Sci U S A.* 1996;93:6830-6834.
- Li RW, Levi DM, Klein SA. Perceptual learning improves efficiency by re-tuning the decision 'template' for position discrimination. *Nat Neurosci.* 2004;7:178-183.
- Zhang JY, Cong LJ, Klein SA, Levi DM, Yu C. Perceptual learning improves adult amblyopic vision through rule-based

- cognitive compensation. *Invest Ophthalmol Vis Sci.* 2014;55:2020-2030.
20. Zhang JY, Zhang GL, Xiao LQ, Klein SA, Levi DM, Yu C. Rule-based learning explains visual perceptual learning and its specificity and transfer. *J Neurosci.* 2010;30:12323-12328.
 21. Zhang JY, Yang YX. Perception learning of motion direction discrimination transfers to an opposite direction with TPE training. *Vision Res.* 2014;99:93-98.
 22. Cong LJ, Wang RJ, Yu C, Zhang JY. Perceptual learning of basic visual features remains task specific with training-plus-exposure (TPE) training. *J Vis.* 2016;16(3):13.
 23. Xiong YZ, Zhang JY, Yu C. Bottom-up and top-down influences at untrained conditions determine perceptual learning specificity and transfer. *eLife.* 2016;5:e14614.
 24. Sengpiel F, Freeman TC, Blakemore C. Interocular suppression in cat striate cortex is not orientation selective. *Neuroreport* 1995;6:2235-2239.
 25. Sengpiel F, Blakemore C, Kind PC, Harrad R. Interocular suppression in the visual cortex of strabismic cats. *J Neurosci.* 1994;14:6855-6871.
 26. Ahissar M, Hochstein S. Attentional control of early perceptual learning. *Proc Natl Acad Sci U S A.* 1993;90:5718-5722.
 27. Pelli DG. The VideoToolbox software for visual psychophysics: transforming numbers into movies. *Spat Vis.* 1997;10:437-442.
 28. Knox PJ, Simmers AJ, Gray LS, Cleary M. An exploratory study: prolonged periods of binocular stimulation can provide an effective treatment for childhood amblyopia. *Invest Ophthalmol Vis Sci.* 2012;53:817-824.
 29. Gao TY, Ledgeway T, Lie AL, et al. Orientation tuning and contrast dependence of continuous flash suppression in amblyopia and normal vision. *Invest Ophthalmol Vis Sci.* 2018;59:5462-5472.
 30. Singer W. The role of attention in developmental plasticity. *Human Neurobiol.* 1982;1:41-43.
 31. Van Balen AT, Henkes HE. Attention and amblyopia an electroencephalographic approach to an ophthalmological problem. *Br J Ophthalmol.* 1962;46:12-20.
 32. Chow A, Giaschi D, Thompson B. Dichoptic attentive motion tracking is biased toward the nonamblyopic eye in strabismic amblyopia. *Invest Ophthalmol Vis Sci.* 2018;59:4572-4580.
 33. Tsirlin I, Colpa L, Goltz HC, Wong AM. Behavioral training as new treatment for adult amblyopia: a meta-analysis and systematic review. *Invest Ophthalmol Vis Sci.* 2015;56:4061-4075.
 34. Levi DM, Klein S. Differences in vernier discrimination for grating between strabismic and anisometric amblyopes. *Invest Ophthalmol Vis Sci.* 1982;23:398-407.
 35. Hess RF, Pointer JS. Differences in the neural basis of human amblyopia: the distribution of the anomaly across the visual field. *Vision Res.* 1985;25:1577-1594.
 36. Gao TY, Anstice N, Babu RJ, et al. Optical treatment of amblyopia in older children and adults is essential prior to enrolment in a clinical trial. *Ophthalmic Physiol Opt.* 2018;38:129-143.
 37. Simonsz-Toth B, Joosse MV, Besch D. Refractive adaptation and efficacy of occlusion therapy in untreated amblyopic patients aged 12 to 40 years. *Graefes Arch Clin Exp Ophthalmol.* 2019;257:379-389.