

Altered frontal connectivity after sleep deprivation predicts sustained attentional impairment: A resting-state functional magnetic resonance imaging study

Ye Cai¹ | Zifeng Mai¹ | Mingzhu Li¹ | Xiaolin Zh

X

1 | INTRODUCTION

Sleep is crucial for survival, but sleep loss commonly occurs in modern society. Individuals experience acute or chronic sleep deprivation due to work pressure, shift work or sleep disorders (Killgore, 2010), and insufficient sleep leads to various dysfunctions of health and

cognitive performance (Lim & Dinges, 2010). Acute sleep deprivation can strongly impair various human cognitive functions, including attention, working memory, executive function, and decision making (Chee et al., 2010; Lim & Dinges, 2008; Xu et al., 2016). Within these cognitive domains, attention is a fundamental component that is very sensitive to sleep deprivation (Killgore, 2010; Lim & Dinges, 2008), which involves

Horne–Ostberg Morningness–Eveningness Questionnaire. The study was conducted in accordance with the Declaration of Helsinki and was approved by the Medical Ethics Committee of South China Normal University. Each participant provided written informed consent; they were paid a certain amount of money after finishing the study.

2.2 | Experimental protocol

The present study was conducted by using a within-subject design. All participants went over two sessions corresponding to rested wakefulness (RW) after normal sleep and acute sleep deprivation, which were counterbalanced by participants and separated by at least 1 week. Prior to the study, participants were asked to attend a semi-structured interview as a screening session to ensure that they

of sleep and habitually awakened between 06:00 and 09:00 hours, as assessed by sleep diary and actigraphy (Actiwatch Spectrum, Philips) within 2 weeks before the in-laboratory experiment. Participants made an additional visit to sleep to the laboratory with polysomnography (PSG) recording 1 night before RW to adapt to the laboratory environment and to ensure that they did not have any sleep disorders. In the RW session, participants had a mean (*SD*) sleep according to the PSG recordings. Resting-state fMRI scanning was performed at 09:30 hours (\pm 1 hr) the next morning. Participants were instructed to lie down and stay awake with their eyes opened. After the scanning session, all participants had to report whether they have fallen asleep in the scanner. One of the participants reported a sleep experience in the scanner and was excluded from data analysis. After the scanning session, participants completed a 10-min PVT. In the sleep-deprivation session, most of the protocol was the same as that in the RW session, and participants went through a 24-hr total sleep-deprivation period during the night, in which participants were

2.3 | Psychomotor vigilance task (PVT) and subjective sleepiness rating

The PVT is widely used to measure sustained attention in sleep research because of its high reliability and high sensitivity to sleep loss (Chua et al., 2019; Lim & Dinges, 2008). In the present study, we

used a 10-min visual PVT. Participants were asked to focus their attention on a red, rectangular box subtending 2×1.3 degrees of visual angle in the middle of a black screen and monitor that space for the appearance of a millisecond counter, which appeared at random intervals ranging from 2 to 10 s. They were instructed to stop the counter as quickly as possible with a button press, after which they would be able to view their RT. Participants were also instructed to avoid anticipating the stimuli so as not to register “false starts” or responses when no stimulus was present on the screen. Trials with RTs of <100 ms were excluded, and those $>$ lapses.

In addition, subjective sleepiness was assessed by the Karolinska Sleepiness Scale (KSS) immediately before the PVT. The KSS is a 9-point, verbally anchored scale ranging from 1 (“extremely alert”) to 9 (“extremely sleepy”).

2.4 | Behavioural data analysis

Behavioural data were processed with SPSS®, version 21 (SPSS Inc.). We used mean RTs and lapse numbers as indices to evaluate individuals’ sustained attention. The change in lapse numbers was calculated as lapse numbers at the sleep-deprivation state – lapse numbers at the RW state. A higher value would represent a more severe sustained attention deficit after sleep deprivation. Two-tailed paired *t* tests were performed to compare lapse numbers between RW and sleep deprivation.

2.5 | Imaging data acquisition

All neuroimaging data were collected under both RW and sleep deprivation conditions with a 3-T Magnetom Trio MRI scanner system (Siemens Medical Systems) using a 12-channel head coil.

echo-planar imaging (EPI) (repetition time [TR] = 2,000 ms, echo time [TE] = 24 ms, slices = 30, slice thickness = 4 mm, matrix size = 64×64 mm², flip angle = 90°, field of view = 220×220 mm², $3.4 \times 3.4 \times 4$ mm³). High-resolution T1-weighted structural imaging was acquired using the magnetisation

TR = 2,300 ms, TE = 3.24 ms, slice thickness = 1 mm, flip angle = 9°,

$$= \quad \times \quad ^2, 1 \times 1 \times 1 \text{ mm}^3$$

were scanned in the axial direction, while T1 structural images were scanned in the sagittal direction.

2.6 | Image pre-processing and statistical analysis

Resting fMRI data pre-processing was performed using the Data Processing Assistant for Resting-State fMRI Advanced Edition

and Analysis of Brain Imaging (DPABI, <http://rfmri.org/DPABI>; Yan et al., 2016) version 4.21 using the following steps: (1) Slice timing and realignment was performed after removing the first 10 time points. (2) Head motion was corrected based on framewise displace-

>0.2 over(r)-1006 BDC 0x0023100426h3E6c4E(020)-B26a00D(T)E31G-1704(S607A0)3000A12A0

TABLE 1 Statistics of behaviour task (PVT) performance and subjective sleepiness rating in both RW and sleep deprivation

Variable, mean (SD)	RW	Sleep deprivation	<i>t</i>
Number of lapses	2.12 (2.66)		
RT, ms			4.69
Subjective sleepiness score			9.60

Abbreviations: PVT, psychomotor vigilance task; RW, rested wakefulness after normal sleep; Mean RT, mean reaction time; subjective sleepiness, a 9-point scale ranging from 1 ("extremely alert") to 9 ("extremely sleepy").

3.2 | Impact of sleep deprivation on ALFF

Paired *t* right paracentral lobule, bilateral thalamus, and inferior occipital gyrus

of the left middle frontal gyrus, right precuneus, right inferior parietal

3.3 | Impact of sleep deprivation on functional connectivity

We performed paired *t* tests to reveal significant differences in the functional connectivity between the RW and sleep-deprivation states. The selected seed regions were based on the regions that - tal region, right paracentral lobule, right thalamus, left middle frontal region, and right inferior parietal lobule. The MNI coordinates and cluster information are provided in Table S1.

3.4 | Correlation between behavioural changes and ALFF/functional connectivity changes

behavioural change

the lapse numbers change ($r = .62, p <$

change and behavioural change

We calculated Pearson's correlation coefficients between the lapse number changes and functional connectivity changes. However, within all the functional connectivity altered by sleep deprivation

(Table S1) only parts of the altered functional connectivity were correlated with the lapse number change. Specifically, the functional connectivity change between the right thalamus and bilateral inferior frontal gyrus was inversely correlated with the lapse number change (right inferior frontal gyrus: $r = p < .001$; left inferior frontal gyrus: $r = p < .001$), while the functional connectivity change between the left middle frontal gyrus and left inferior occipital gyrus was significantly correlated with the lapse number change ($r = .68, p <$

4 | DISCUSSION

In the present study, we investigated the effect of sleep deprivation

explored the neural basis of vigilance impairment after acute sleep deprivation by connecting changes in PVT performance with altered functional connectivity in resting-state fMRI data. The present results demonstrated a widespread detrimental effect of sleep deprivation

ual and motor regions were significantly increased, which may suggest a counteracting effect to keep a person alert when combating

were correlated with the lapse numbers change. Critically, correlation analyses revealed that a decrease in the functional connectivity between the right thalamus and bilateral inferior frontal gyrus was significantly correlated with increased lapses during the PVT, while an increase in the functional connectivity between the left middle frontal gyrus and left inferior occipital gyrus was significantly correlated with increased lapses during PVT. These findings indicated that decreased co-activation of the thalamus and bilateral inferior frontal gyrus might predict vigilance impairment after sleep deprivation, as well as increased co-activation between frontal and visual areas.

ent study, including the superior frontal gyrus and right precuneus,

This is consistent with our assumption. Several studies concern-

connectivity was disrupted in a sleep-deprived state (De Havas et al., 2012; Gujar et al., 2010). These results jointly support our present results that sleep deprivation disrupts resting DMN function, which might be related with reduced responsiveness to the environment. Consistent with our hypothesis, the right inferior parietal lobule and right superior frontal gyrus were found to ex-

with attention and working memory (Corbetta & Shulman, 2002;

mus, visual cortex, and paracentral lobule after sleep deprivation was consistent with several recent studies and meta-analysis (Gao

the thalamus after sleep deprivation might represent a compensatory mechanism to maintain relative alertness, as mentioned in a

visual cortex and motor cortex might correspond to the demand to enhance perceptual function to keep the person alert and aware of

the attention-related regions, sleep deprivation has led to poorer attention function, and thus, increased perceptual load of visual pro-

the visual cortex and the lapse number change in the present study.

value in the visual cortex, which might indicate that the hyperactivity in several brain regions found in the present study might be interpreted as an enhanced neural function to confront the impact caused by sleep deprivation.


Correlation analysis revealed that the functional connectivity between the thalamus and bilateral inferior frontal gyrus is correlated with the vigilance decline induced by sleep deprivation. The disruption of thalamic connectivity with frontal regions might reflect “microsleeps”, which indicate instability of the waking state. A mi-

into wakefulness, especially when individuals are sleep deprived


(Goel et al., 2009). Previous studies have observed transient re-

occurred. The frequently inconsistent altered activity in the frontal cortex and thalamus might parallel decreased synchronisation in the thalamo-cortical connectivity (Poudel et al., 2018; Shao et al., 2013). Consistent with these studies, the present results demonstrated that participants who had reduced thalamo-cortical connectivity had more lapses in the subsequent PVT, which indicated a stronger unstable state and might reflect the neural basis of individual differences of vulnerability to sleep deprivation. In addition, a review demonstrated that as an important node of the arousal system, the thalamus is part of the stimulus-driven control network, anatomically connecting with the temporoparietal junction and inferior frontal cortex (Corbetta & Shulman, 2002). Therefore, the reduced frontal connectivity with the thalamus seen in the present study might be


(a)




Seed Region: R. Thalamus
(9, -18, 6)




(b)




Seed Region: R. Thalamus
(9, -18, 6)



(c)



Seed Region: L. MFG
(-27, 6, 45)



intrinsic brain activity alterations in unmedicated depression. *Journal of Affective Disorders*, 172

fMRI signals predicts Stop signal task performance. *NeuroImage*, 60,

default mode network: Inter-subject variability and effects of sleep deprivation. *Brain Research*, 1596

The study on functional connectivity between frontal eye field and other brain cortex in visual processing. Paper presented at the 2010 International Conference on Computational Aspects of Social Networks, 2010:333–336. <https://doi.org/10.1109/CASoN.2010.81>

individual differences in vulnerability toward total sleep deprivation-induced changes in cognitive function. *Journal of Sleep Research*, 25

cessing analysis for (resting-state) brain imaging. *Neuroinformatics*, 14