

Crowding alters the spatial distribution of attention modulation in human primary visual cortex

Department of Psychology and Key Laboratory of Machine Perception (Ministry of Education), Peking University, Beijing, PR China

Fang Fang



Sheng He

Department of Psychology, University of Minnesota, Minneapolis MN, USA



Crowding effect is the visibility reduction of a target when presented with neighboring distractors. It has been explained by either lateral inhibition at a pre-attentive level or coarse spatial resolution of attention. To test these theories, high-resolution fMRI was used to measure V1 response to the target in the presence or the absence of the distractors in both attended and unattended conditions. We found the cortical response to the target was not affected by the presence of distractors in the unattended condition. However, the spatial distribution of attention modulation in the target and its surrounding area depended on the crowding configuration. When distractors were placed in the same radial axis as the target, a configuration with a severe crowding effect, significant attention enhancements were observed not only in the target's and the distractors' locations, but also in regions next to the target where even no stimulus was presented. But this spread of attention enhancement did not occur when distractors were placed in the same circumference as the target, a configuration with a weak crowding effect. The pattern of interaction between attention and target-distractor configuration supports that crowding results from coarse spatial resolution of attention.

Keywords: attention, contrast, crowding, fMRI, vision, V1

Citation: Fang, F., & He, S. (2008). Crowding alters the spatial distribution of attention modulation in human primary visual cortex. *Journal of Vision*, 8(9):6, 1–9, <http://journalofvision.org/8/9/6/>, doi:10.1167/8.9.6.

Introduction



(Fang & He, 2008). (Fang, 1A). (B... 1970; ... 1991). (B... 1970; ... 1991). (C... 1997). (L... & L... 1997). (L... K... & ... 1985; ... & E... 1986). (L... & ... 1988). (S... & M... 1976). (L... & E... 1997). (L... & M... 2004). (L... B... & ... 2007; M... M... & ... 2005). (B... L... & ... 1979; L... & ... 1987). (L... 2004). (L... & ... 1973; ... & ... 1983). (H... C... & I... 1996, 1997; I... & C... 2001). (S... 2005; S... 1991). (C... 1991). (B... 1970; S... & B... 1962). (M... I: (1) H... (2) H... ?

(S... 2001). (S... 2005; S... 1991). (C... 1991). (B... 1970; S... & B... 1962). (M... I: (1) H... (2) H... ?

Methods

Participants

A... (3... 4...) fMRI



MRI data acquisition

The subjects were scanned using a 3T MRI scanner (60 H). The functional MRI (fMRI) data were acquired using a BOLD (Blood Oxygen Level Dependent) sequence. The parameters were: TR = 2000 ms, TE = 35 ms, FOV = 22 × 22 cm², matrix = 128 × 128, slice thickness = 75 mm, and 20 slices. The anatomical MRI (M1) was acquired using a 3D T1-weighted sequence (3D MAGE; 1 × 1 × 1 mm). The functional MRI (fMRI) data were acquired using a BOLD sequence with the following parameters: TR = 2000 ms, TE = 35 ms, FOV = 22 × 22 cm², matrix = 128 × 128, slice thickness = 75 mm, and 20 slices. The anatomical MRI (M1) was acquired using a 3D T1-weighted sequence (3D MAGE; 1 × 1 × 1 mm). The functional MRI (fMRI) data were acquired using a BOLD sequence with the following parameters: TR = 2000 ms, TE = 35 ms, FOV = 22 × 22 cm², matrix = 128 × 128, slice thickness = 75 mm, and 20 slices.

MRI data processing and analysis

The functional MRI (fMRI) data were processed using SPM99 (Statistical Parametric Mapping) software. The anatomical MRI (M1) was processed using the same software. The functional MRI (fMRI) data were processed using the following parameters: TR = 2000 ms, TE = 35 ms, FOV = 22 × 22 cm², matrix = 128 × 128, slice thickness = 75 mm, and 20 slices. The anatomical MRI (M1) was processed using the following parameters: TR = 2000 ms, TE = 35 ms, FOV = 22 × 22 cm², matrix = 128 × 128, slice thickness = 75 mm, and 20 slices.




Figure 2. Regions of interest. (A) Five flickering round checked patches with a full contrast were used to define the ROIs (central, upper, lower, left and right). They occupied the same spatial extents as the target and the distractors. (B) Cortical activations by the five patches are depicted in a representative inflated brain. The red, green, blue, yellow, and light blue areas correspond to the left, central, right, lower, and upper ROIs, respectively. V1 is defined by retinotopic mapping and its boundaries are indicated by the white dashed lines.




Figure 3. Behavioral and cortical responses to the target in the single, tangential and radial configurations. (A) Performance in the contrast discrimination task. (B) BOLD responses to the target with the luminance discrimination task at the fixation point. Error bars denote 1 SEM calculated across subjects.

Eye movement recording

Eye movement recording was performed using a 60 Hz video camera (A, 3) and a video camera (A, 3) to record eye movements. The eye movement recording was performed using a video camera (A, 3) and a video camera (A, 3) to record eye movements. The eye movement recording was performed using a video camera (A, 3) and a video camera (A, 3) to record eye movements.

Results

Behavioral performance was significantly higher in the radial configuration compared to the tangential configuration ($F_{(1,14)} = 17.962, p < 0.001$),




Figure 3. Behavioral and cortical responses to the target in the single, tangential and radial configurations. (A) Performance in the contrast discrimination task. (B) BOLD responses to the target with the luminance discrimination task at the fixation point. Error bars denote 1 SEM calculated across subjects.




Figure 4. BOLD signals in the left, right, upper, lower, and central ROIs in the single, tangential, and radial configurations when subject attended to either the fixation (left part of a panel) or the target (right part of a panel). Error bars denote 1 SEM calculated across subjects.


(C & S, 2002). I
 OI
 C
 (t = 3.118, p = 0.036) (t = 3.623, p = 0.022) OI,
 OI (F, 5A,
 (t = 3.64, p = 0.022) (t = 3.863, p = 0.018) OI (F, 5A,
 (F, 4B).
 E (F, 6
 (M I

... ($t < 2$, $p > 0.12$).

Discussion

... (...-L ... & H ... , 2003).
H ... , ... (... , ... , ...)
... -L ... H ...
... (M I ... (... , ... - ...) . I ... , A ... ,
C ... (2006) .
...
... I ...
...) 8. 282-298 (...)--414 (8)- M ... 8

...
... M...
... € ... €
... € ... €



References

Abrams, G. J., & Kornhuber, H. H. (1963). Localization of the body in space. *Perception & Psychophysics*, 53, 658–667. [M](#)

Abrams, A. C., Cornblath, S. L., & Mountcastle, V. B. (2006). Neural basis of body representation. *Journal of Vision*, 6(6):804, 804. [M](#)

Baltes, P. B., Lindenberger, D., & Staudt, M. (1979). The aging of human intelligence. *Perception & Psychophysics*, 25, 447–456. [M](#)

Baltes, P. B. (1970). Intelligence. *Nature*, 226, 177–178. [M](#)

Caspari, J., Bressan, J. P., & Mountcastle, V. B. (2002). Neural basis of body representation. *Journal of Neurophysiology*, 88, 2530–2546. [M](#)

Caspari, M., & Mountcastle, V. B. (2002). Cortical representation of body representation. *Nature Reviews, Neuroscience*, 3, 201–215. [M](#)

Davatzikos, C., & Mountcastle, V. B. (2003). Cortical representation of body representation. *Neuron*, 38, 659–671. [M](#)

Eickhoff, S. B., Grafton, G. H., & Hallett, H. (1997). The representation of body representation. *Cerebral Cortex*, 7, 181–192. [M](#)

Haxby, J. V., Gauthier, I., & Tarr, P. J. (1996). A cortical representation of body representation. *Nature*, 383, 334–337. [M](#)

Haxby, J. V., Gauthier, I., & Tarr, P. J. (1997). A cortical representation of body representation. *Trends in Cognitive Sciences*, 1, 115–121. [M](#)

Iacoboni, M., & Grafton, G. H. (2001). Cortical representation of body representation. *Cognitive Psychology*, 43, 171–216. [M](#)

Lewis, D., & Mountcastle, V. B. (1989). Cortical representation of body representation. *Psychological Review*, 96, 101–124. [M](#)

Lewis, D. M., Kornhuber, H. H., & Mountcastle, V. B. (1985). Cortical representation of body representation. *Vision Research*, 25, 963–977. [M](#)

Lewis, D. M., Kornhuber, H. H., & Mountcastle, V. B. (1987). Cortical representation of body representation. *Vision Research*, 27, 581–597. [M](#)

Lewis, J. B., & Mountcastle, V. B. (1997). Cortical representation of body representation. *Nature*, 387, 73–76. [M](#)

Lewis, E. G., Bressan, J. P., & Mountcastle, V. B. (2007). Cortical representation of body representation. *Journal of Vision*, 7(2):24, 1–11. [M](#)

Mountcastle, V. B., Mountcastle, N. J., & DeYoe, E. A. (2005). A cortical representation of body representation. *Journal of Vision*, 5(1):6, 58–70. [M](#)

Mountcastle, V. B., Mountcastle, N. J., & DeYoe, E. A. (2004). Cortical representation of body representation. *Journal of Vision*, 4(12):12, 1136–1169. [M](#)

Mountcastle, V. B., Mountcastle, N. J., DeYoe, E. A., & Bressan, J. P. (1995). Body representation. *Science*, 268, 889–893. [M](#)

Mountcastle, V. B., Mountcastle, N. J., DeYoe, E. A., & Bressan, J. P. (1999). Cortical representation of body representation. *Neuroimage*, 9, 526–533. [M](#)

Mountcastle, V. B. (2005). Cortical representation of body representation. *Journal of Vision*, 5(11):8, 1024–1037. [M](#)

Mountcastle, V. B., Mountcastle, N. J., & DeYoe, E. A. (1991). Cortical representation of body representation. *Perception & Psychophysics*, 49, 495–508. [M](#)

Mountcastle, V. B., & Mountcastle, N. J. (1962). A cortical representation of body representation. *American Journal of Ophthalmology*, 53, 471–477. [M](#)

Mountcastle, V. B., & Mountcastle, N. J. (1992). Cortical representation of body representation. *Vision Research*, 32, 1349–1357. [M](#)

Mountcastle, V. B., & Mountcastle, N. J. (1973). Cortical representation of body representation. *Psychological Review*, 80, 284–302. [M](#)

Mountcastle, V. B., Mountcastle, N. J., & Mountcastle, N. J. (1976). Cortical representation of body representation. *Journal of the Optical Society of America A*, 66, 332–338. [M](#)

Mountcastle, V. B., & Mountcastle, N. J. (1988). Cortical representation of body representation. *American Journal of Optometry and Physiological Optics*, 65, 395–399. [M](#)

Wang, F., Zhang, H., & Egeton, D. (1997). Lateral masking of visual motion. *Journal of the Optical Society of America A*, *14*, 2057–2068. [M](#)

Wang, F. A., & Egeton, E. A. (1986). A study of visual motion. *Vision Research*, *26*, 349–360. [M](#)

Wang, G., & Chang, L. (1983). Lateral masking of visual motion. *Perception & Psychophysics*, *33*, 129–138. [M](#)

Wang-Li, B., & He, D. J. (2003). Lateral masking of visual motion. *Journal of Neuroscience*, *23*, 6884–6893. [M](#) [A](#)